

# Finite Elements Homework 1

Lisandro Agustin Roldan

November 10, 2015

## PROBLEM

Consider the following differential equation

$$-u'' = f \text{ in } [0, 1]$$

with the boundary conditions  $u(0) = 0$  and  $u(1) = \alpha$ .

The Finite Element discretization is a 2-noded linear mesh given by the nodes  $x_i = ih$  for  $i = 0, 1, \dots, n$  and  $h = 1/n$ .

1. Find the weak form of the problem. Describe the FE approximation  $u^h$ .
2. Describe the linear system of equations to be solved.
3. Compute de FE approximation  $u^h$  for  $n = 3$ ,  $f(x) = \sin(x)$  and  $\alpha = 3$ . Compare it with the exact solution,  $u(x) = \sin(x) + [3 - \sin(x)]x$ .

## SOLUTION

The strong form of the problem is given by:

$$A(u) = \frac{d^2u}{dx^2} + Q = 0 \text{ in } [0, 1]$$

$$B(u) = u - \bar{u} = 0 \text{ in } x = 0 \text{ and } x = 1$$

We don't have any Neumann boundary condition, therefore the integral form of the problem is:

$$\int_0^1 \omega \left[ \frac{d^2u}{dx^2} + Q \right] dx = \int_0^1 \omega \frac{d^2u}{dx^2} dx + \int_0^1 \omega Q(x) dx = 0$$

Integrating by parts the first term of the equation we have

$$- \int_0^1 \frac{d\omega}{dx} \frac{du}{dx} dx + \omega \frac{du}{dx} \Big|_0^1 + \int_0^1 \omega Q(x) dx = 0$$

We can approximate the function  $u(x)$  as  $u^h(x) = N_i(x)u_i$ , call  $\frac{du}{dx}$  the reaction force/flux  $q$ , use the Galerking method choosing  $\omega(x) = N_i(x)$  and rearrange the terms of the equation. Then, the weak form of the problem will be:

$$\int_0^1 \frac{dN_i}{dx} \sum_{j=1}^n \left( \frac{dN_j}{dx} u_j \right) dx = N_i q \Big|_0^1 + \int_0^1 N_i Q(x) dx$$

This expression represents a linear system of equation of the form  $K_{ij} u_j = f_i$ .

$K_{ij}$  is a square symmetric matrix of size  $n \times n$ . Where  $n$  is the number of nodes in the discretization.

$$K_{ij} = \int_0^1 \frac{dN_i}{dx} \frac{dN_j}{dx} dx$$

$u_j$  is a vector of size  $n$  with the values of the unknown in the nodes.

$f_i$  is a vector of size  $n$  with the independent terms of the equation.

$$f_i = N_i q \Big|_0^1 + \int_0^1 N_i Q(x) dx$$

$$\begin{bmatrix} K_{11} & K_{12} & K_{13} & \dots & K_{1n} \\ K_{21} & K_{22} & K_{23} & \dots & K_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ K_{n1} & K_{n2} & K_{n3} & \dots & K_{nn} \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \dots \\ u_n \end{bmatrix} = \begin{bmatrix} f_1 \\ f_2 \\ \dots \\ f_n \end{bmatrix}$$

For a 3 finite element discretization of the domain we can find the system of equation to solve by assembling the local components of  $K$  and  $f$  of each element.

For each element, the  $K$  matrix will be

$$K^e = \frac{1}{1/3} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$$

And the  $f$  vector will depend on the integration over the length of the element of the function  $Q(x) = \sin(x)$  plus the boundary flux/reaction  $q$  when pertinent.

Knowing that the locally the shape function has the form of:

$$N_1^e = \frac{x_2 - x}{l^e}$$

$$N_2^e = \frac{x - x_1}{l^e}$$

We can find the 6 shape functions  $N_i$  for each node of the 3 elements.

$$N_1^1 = 3(1/3 - x)$$

$$N_2^1 = 3(x - 0)$$

$$N_1^2 = 3(2/3 - x)$$

$$N_2^2 = 3(x - 1/3)$$

$$N_1^3 = 3(1 - x)$$

$$N_2^3 = 3(x - 2/3)$$

$f$  will take a value in each node of the elements of:

$$f_1^1 = \int_0^{1/3} (-3x + 1) \sin(x) dx + q_1 = 0.018 + q_1$$

$$f_2^1 = \int_0^{1/3} (3x) \sin(x) dx = 0.037$$

$$f_1^2 = \int_{1/3}^{2/3} (-3x + 2) \sin(x) dx = 0.071$$

$$f_2^2 = \int_{1/3}^{2/3} (3x - 1) \sin(x) dx = 0.088$$

$$f_1^3 = \int_{2/3}^1 (-3x + 3) \sin(x) dx = 0.117$$

$$f_2^3 = \int_{2/3}^1 (3x - 2) \sin(x) dx + q_4 = 0.129 + q_4$$

Assembled, the general system of equation will take the form of:

$$\begin{bmatrix} K_{11}^1 & K_{12}^1 & 0 & 0 \\ K_{21}^1 & K_{22}^1 + K_{11}^2 & K_{12}^2 & 0 \\ 0 & K_{21}^2 & K_{22}^2 + K_{11}^3 & K_{12}^3 \\ 0 & 0 & K_{21}^3 & K_{22}^3 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{bmatrix} = \begin{bmatrix} f_1^1 \\ f_2^1 + f_1^2 \\ f_2^2 + f_1^3 \\ f_2^3 \end{bmatrix}$$

Replacing the values of  $K$  and  $f$  previously found and introducing the Dirichlet boundary conditions we get ( $\alpha = 3$ ):

$$3 \begin{bmatrix} 1 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ u_2 \\ u_3 \\ 3 \end{bmatrix} = \begin{bmatrix} 0.018 + q_1 \\ 0.108 \\ 0.204 \\ 0.129 + q_4 \end{bmatrix}$$

As  $u_1$  and  $u_4$  are known values we can reduce the system to:

$$3 \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} u_2 \\ u_3 \end{bmatrix} = \begin{bmatrix} 0.108 \\ 0.204 + 9 \end{bmatrix}$$

Solving the system we find that  $u_2 = 1.059$  and  $u_3 = 2.051$


Now, with all the node values known we can compute the reactions/fluxes  $q_1$  and  $q_4$

$$-1.059 = 0.018 + q_1 \rightarrow q_1 = -1.007$$

$$-2.051 + 3 = 0.129 + q_4 \rightarrow q_4 = -0.820$$

We also can compute the values of  $u$  in the same points but using the real governing equation  $u(x) = \sin(x) + [3 - \sin(x)]x$

Node	$u_{fem}$	$u_{exact}$	Difference
1	0	0	0
2	1.059	1.046	0.013
3	2.051	2.057	-0.006
4	3	3	0



*The red line represents the exact solution while the blue points represent the values found with the finite element method.*

The approximation was expected to be slightly different from the exact solution because of the finite method itself and the computational tools used to compute the integrals and solve the system of equation.

Finally, we can compute the mean squared error applying the following expression:

$$MSE_{Error} = \frac{1}{n} \sum_{i=1}^n (u_{fem} - u_{exact})^2 \approx 5 \cdot 10^{-5}$$